

Impairment in personality functioning predicts young adult suicidal ideation and suicide attempt above and beyond depressive symptoms

Kiran Boone, Kennedy M. Balzen, Carla Sharp

doi: 10.4081/ripppo.2024.814

Supplementary Material

Supplementary Table 1. *Descriptive Statistics of Study Variables in Clinical Sample (N=63).*

	Minimum	Maximum	Mean	Std. dev.	Skewness	Kurtosis
Age	18	25	22.22	2.20	-0.54	-0.84
Suicidal ideation	3.00	38.00	20.64	8.69	0.19	-0.80
Number of suicide attempts	0	11	2.06	3.04	1.74	2.07
Overall personality functioning	16.00	46.00	34.75	6.33	-0.57	0.21
Self-functioning	8.00	24.00	18.92	3.50	-0.73	0.51
Interpersonal functioning	8.00	24.00	15.83	3.73	-0.03	-0.47
Depression symptoms	0.00	3.92	2.11	1.01	-0.12	-0.67

Std. dev., standard deviation.

Supplementary Table 2. *Descriptive Statistics of Study Variables in College Sample (N=90).*

	Minimum	Maximum	Mean	Std. dev.	Skewness	Kurtosis
Age	18	24	20.03	1.61	0.75	0.01
Suicidal ideation	2.00	30.00	8.27	5.62	1.24	1.91
Number of suicide attempts	0	6	0.20	0.94	5.59	32.29
Overall personality functioning	12.00	40.00	24.70	6.55	-0.16	-0.57
Self-functioning	6.00	22.00	13.44	3.96	-0.23	-0.71
Interpersonal functioning	6.00	20.00	11.26	3.43	0.32	-0.62
Depression symptoms	0.00	3.09	1.20	0.92	0.49	-1.09

Std. dev., standard deviation.